

1. Consider the following propositional theory

$$T_n = \{r_1 \rightarrow r_2, r_2 \rightarrow r_3, \dots, r_{n-1} \rightarrow r_n, r_n \rightarrow r_1\}$$

based on $\mathcal{P}_n = \{r_1, \dots, r_n\}$ where $n > 0$. The intuitive reading of r_i is that the i^{th} node is *reachable* in a finite ring of n nodes.

- (a) Determine the models $M \subseteq \mathcal{P}_n$ of T_n .
 - (b) Show/argue that $T \models r_i \rightarrow r_j$ for any $0 < i, j \leq n$.
 - (c) Compare the models of T_n with the models of $T_n \cup \{r_1\}$? Does T_n properly formalize reachability among the nodes in the ring?
2. Consider a set of nodes $U_n = \{1, \dots, n\}$ and a relation $E \subseteq (U_n)^2$ representing the edge relation of a graph $G = \langle U_n, E \rangle$ with n vertices. Which properties of G are captured by the following first-order sentences?

- (a) $\phi_1 = \forall x \exists y E(x, y) \wedge \forall x \forall y \forall z (E(x, y) \wedge E(x, z) \rightarrow y = z)$
- (b) $\phi_2 = \forall x \exists y E(y, x) \wedge \forall x \forall y \forall z (E(y, x) \wedge E(z, x) \rightarrow y = z)$
- (c) $\phi_3 = \phi_1 \wedge \phi_2$
- (d) $\phi_4 = \forall x E(x, x)$
- (e) $\phi_5 = \forall x \forall y (E(x, y) \rightarrow E(y, x))$
- (f) $\phi_6 = \forall x \forall y \forall z (E(x, z) \wedge E(z, y) \rightarrow E(x, y))$
- (g) $\phi_7 = \phi_4 \wedge \phi_5 \wedge \phi_6$

3. Describe the models of ϕ_3 and ϕ_7 (see above) when $n = 3$.

4. Consider the following first-order theory

$$T = \{N(0, s(0)), \forall x \forall y (N(x, y) \rightarrow N(s(x), s(y)))\}$$

which captures for each natural number its immediate successor.

- (a) Determine the Herbrand base $\text{Hb}(T)$.
- (b) Determine a Herbrand model $M \subseteq \text{Hb}(T)$ for T .
- (c) Is M minimal with respect to \subseteq , i.e., is there another Herbrand interpretation $M' \subset M$ of T such that $M' \models T$?