

T-79.5201 Discrete Structures, Autumn 2006

Tutorial 7, 22 November

1. Verify the following properties of the Riemann–Stieltjes -integral, assuming that the integrals appearing in the formulas are well defined:

(a) Linearity:

$$\begin{aligned}\int_a^b (c_1 f_1 + c_2 f_2) dg &= c_1 \int_a^b f_1 dg + c_2 \int_a^b f_2 dg, \\ \int_a^b f d(c_1 g_1 + c_2 g_2) &= c_1 \int_a^b f dg_1 + c_2 \int_a^b f dg_2.\end{aligned}$$

(b) Reduction to Riemann integral: for a continuously differentiable function g ,

$$\int_a^b f(t) dg(t) = \int_a^b f(t)g'(t) dt.$$

2. Let f and g be continuous functions and $a, b \in \mathbf{Z}$. Verify the correctness of the following formulas, assuming that the integrals contained in them are well defined:

$$\begin{aligned}\int_a^b f(t) dg(\lceil t \rceil) &= \sum_{a \leq k < b} f(k) \Delta g(k), & \Delta g(k) &= g(k+1) - g(k); \\ \int_a^b f(\lceil t \rceil) dg(t) &= \sum_{a < k \leq b} f(k) \nabla g(k), & \nabla g(k) &= g(k) - g(k-1).\end{aligned}$$

Derive from the preceding formulas the following “partial summation rule”:

$$\sum_{a \leq k < b} f(k) \Delta g(k) = \int_a^b f(k)g(k) - \sum_{a < k \leq b} g(k) \nabla f(k).$$

3. Use Euler’s summation formula to estimate the following sums:

- (a) Sum $\sum_{1 \leq k < n} k^{1/2}$ up to order $O(1)$.
- (b) Sum $\sum_{1 \leq k < n} k^r$, $r \in \mathbf{N}$, exactly.